



COMPOUNDED INJECTABLE TIRZEPATIDE

Patient _____

Prescriber _____

DOB _____

Phone _____

Phone _____

Fax _____

Address _____

Address _____

Compounded Tirzepatide Injections

Vial: 10mg/mL

Instructions/Usual Starting Dose:

Inject 0.25mL (2.5mg) weekly subcutaneously for first 4 weeks.

Inject 0.5mL (5mg) weekly subcutaneously for weeks 5 -8

Taper up as tolerated and needed.

*FDA recommends disposing of vial 28 days after 1st puncture.

☐ **Vial: 20mg/2mL**

SIG: _____

Refill _____ times

Cash Price: \$300

Includes 5 syringes, alcohol pads, tirzepatide vial, mailing

☐ **Ondansetron 4mg ODT**

SIG: Place one tablet on tongue and allow to dissolve every 4-6 hours as needed for nausea. Take first tablet 30 minutes prior to semaglutide dose increase.

Cash Price: \$10.00

10 Tablets

Refill _____ times

Mailing Options

☐ Send compound to patient, bill the patient |

☐ Send compound to clinic

☐ Send compound patient, bill the clinic

Rev:4/30/2024

SIGNATURE _____

Date _____

City Drug, 131 10th St, Evanston (307) 789-4000